

# The dry matter yield and nutritive value of wet-tolerant tropical forage legumes in single cropping or mixed cropping with gramineous forage crops in drained paddy field

M. TOBISA<sup>1</sup>, Y. NAKANO<sup>2</sup>, K. OKANO<sup>2</sup>, M. SHIMOJO<sup>2</sup> AND Y. MASUDA<sup>2</sup>

<sup>1</sup> Faculty of Agriculture, University of Miyazaki, Miyazaki, 889-2192, Japan,

E-mail: mtobisa@cc.miyazaki-u.ac.jp,

<sup>2</sup> Faculty of Agriculture, Kyushu University, Fukuoka, 812-8581, Japan

## Introduction

In Japan, the production of rice has been controlled since the 1970s and some parts of the paddy fields have been laid off for forage production. However, in poorly-drained fields or fields with high ground water table, forage species with high tolerance of wet conditions are required. The tropical forage legumes *Aeschynomene americana* cv. Glenn (Glenn) and *Macroptilium lathyroides* cv. Murray (phasey bean) have a high wet endurance (Bishop *et al.* 1985; Tobisa *et al.* 1999) and show high dry matter productivity (Skerman *et al.* 1988; Tobisa *et al.* 1999). The objective of this experiment was to evaluate the dry matter yield and nutritive value of Glenn and phasey bean in single cropping or mixed cropping with gramineous forage crops in drained paddy fields.

## Materials and methods


The experiment was conducted in a drained paddy field adjoining a rice paddy field at the Kyushu University Farm. Tropical forage legumes (Glenn and phasey bean), gramineous forage crops [Japanese banyard millet (cv. White panic and Aoba millet), maize (cv. Snow dent 123) and sorghum (cv. Ultra sorgo)] were used. On 23 June, the paddy field was sown at 2 kg/ha in rows with a distance of 50 cm between rows with alternate row sowings of the legume and the gramineous forage crop. Plants were harvested on September 2 and 22 and November 18 (data not shown), and measurements made for dry matter yield (DMY), *in vitro* dry matter digestibility (IVDMD) (Goto and Minson 1977) and crude protein (CP). Digestible dry matter yield (DDMY) and CP yield (CPY) were calculated.

## Results

Dry matter yield of the single-cropped Glenn was similar to that of the single-cropped sorghum when cut on September 22 and November 18. The DDMY of Glenn-sorghum mixed crop was higher than that of the single-cropped sorghum when cut on September 22. The mixed crops of forage legumes and gramineous forages showed higher total CPY compared with the single crop of gramineous forage when cut on September 22.

## Conclusions

The results of the present study demonstrated that Glenn has a high DMY, DDMY and CPY. The mixed crop of forage legume with a gramineous forage with high wet tolerance provided a good forage production system for the drained paddy field.



G: Glenn, P: Phasey bean, W: White panic,

A: Aoba millet, M: Maize, S: Sorghum.

The values with different letters are significantly different at  $P < 0.05$  at each sampling time.

■ Legume □ Gramineous forage

**Figure 1.** Dry matter yield (DMY, 1), digestible dry matter yield (DDMY, 2) and crude protein yield (CPY, 3) of single and mixed crops in the drained paddy field.

## References

- BISHOP, H.G., LUDKE, D.H. and RUTHERFORD, M.T. (1985) Glenn jointvetch: A new pasture legume for Queensland coastal areas. *Queensland Agricultural Journal*, **111**, 241–245.
- GOTO, I. and MINSON, D.J. (1977) Prediction of the dry matter digestibility of tropical grasses using a pepsin-cellulase assay. *Animal Feed Science and Technology*, **2**, 247–253.
- SKERMAN, P.J., CAMERON, D.G. and RIVEROS, F. (1988) In: *Tropical Forage Legumes*. pp. 205–211. (FAO: Rome).
- TOBISA, M., SHIMOJO, M., OKANO, K. and MASUDA, Y. (1999) Growth habit of tropical forage legume genus *Aeschynomene* in the drained paddy field and upland field. *Grassland Science*, **45**, 248–256. (In Japanese).